

## ORIGINAL ARTICLE

## *Listeria monocytogenes* Contamination in Unpasteurized Traditional Cheese Products in Qazvin, Iran

Hajar Khedmati Morasa<sup>1</sup>,  
Razzagh Mahmoudi<sup>2</sup>,  
Peyman Ghajarbeygi<sup>3</sup>,  
Shaghayegh Mosavi<sup>4</sup>,  
Saeed Shahsavari<sup>5</sup>,  
Narges Abbasi<sup>1</sup>,  
Neda Sarfalah<sup>1</sup>

<sup>1</sup> MSc in Food Safety and Health, Department of Food Safety and Health, School of Public Health, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>2</sup> Professors, Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>3</sup> Associate Professor, Health Products Safety Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>4</sup> PhD Student in Molecular Medicine, Faculty of Medical Sciences, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>5</sup> Instructor in Biostatistics, Health Products Safety Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

(Received April 22, 2019 ; Accepted August 25, 2019)

### Abstract

**Background and purpose:** *Listeria monocytogenes* is a foodborne pathogen and a potential risk to public health. Listeriosis is one of the most serious infectious diseases in most developed countries. Consumption of raw milk and unpasteurized traditional dairy products such as cheese can be a major reason for listeriosis in humans. This research aimed at investigating *Listeria monocytogenes* contamination in unpasteurized cheese products by using culture and Polymerase Chain Reaction (PCR) in Qazvin, Iran.

**Materials and methods:** In this research, 128 samples of traditional cheese products were collected from different traditional shopping centers in Qazvin, between October 2017 and September 2018. They were transported to the laboratory under controlled conditions. All isolates were analysed to biochemical test. *L. monocytogenes* strains were further confirmed by PCR amplification.

**Results:** Findings showed that 14 samples (10.9%) were contaminated with *L. monocytogenes*. The highest prevalence of *L. monocytogenes* was found in white cheese samples (7%). The highest rate of contamination was reported in spring and winter (3.1%).

**Conclusion:** *Listeria* contamination in cheese samples studied can pose a serious risk to consumers of non-pasteurized dairy products. Therefore, food safety and health practitioners should apply effective methods and standards.

**Keywords:** *Listeria monocytogenes*, unpasteurized cheese, Qazvin, PCR

J Mazandaran Univ Med Sci 2019; 29 (178): 115-126 (Persian).

\* Corresponding Author: Razzagh Mahmoudi- Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran (E-mail: r.mahmoudi@yahoo.com)